arXiv:gr-qc/0306073v2 18 Jun 2003

An explicit global covering of the Schwarzschild-Tangherlini black holes

Kayll Lake [*]
Department of Physics and Department of Mathematics and Statistics,
Queen's University, Kingston, Ontario, Canada, K7L 3N6
(Dated: February 7, 2008)

Explicit regular coordinates that cover all of the Tangherlini solutions (Schwarzschild black holes of dimension $D > 4$) are given. The coordinates reduce to Israel coordinates for $D = 4$.

One consequence of string theory is the importance now attached to the study of higher dimensional black hole backgrounds. Even at five dimensions asymptotically flat stationary vacuum black holes are not unique in the sense that horizons of topology $S^1 \times S^2$ [1] in addition to S^3 [2] are now known. However, in the simpler static case, it is known that asymptotically flat static vacuum black holes are unique [3] and given by the Tangherlini generalization of the Schwarzschild vacuum [4]. The purpose of this note is to exhibit global, regular and explicit coordinates for these spaces. In addition to their obvious utility, these coordinates may be useful for studies of stability in the nonlinear regime [5].

For dimensions $D \geq 4$ the space

$$ds^2 = -2\,\frac{w^2(-(r)^{-D+3}2^D m^D + 4m^2(D(2m-r)+3r-4m)}{m(2\,m-r)^2}dU^2 + 8m dU dw + r^2 d\Omega_{D-2}^2, \tag{1}$$

where $r \geq 0$ signifies the function

$$r(U,w) \equiv 2\,m + (D-3)Uw, \tag{2}$$

m is a constant > 0 and $d\Omega_{D-2}^2$ is the metric of a unit $D-2$ sphere [6], satisfies

$$R_{ab} = 0 \tag{3}$$

where R_{ab} is the Ricci tensor [7]. The trajectories with D-tangents $k^a = \delta_w^a$ are null geodesics affinely parameterized by w and give $r\nabla_a k^a = (D-2)(D-3)U$. For (1) there is but one independent invariant derivable from the Riemmann tensor without differentiation. This invariant can be taken to be the square of the Weyl tensor C_{abcd} and for (1)

$$C_{abcd}C^{abcd} \propto \frac{m^{2(D-3)}}{r^{2(D-1)}} \tag{4}$$

and so (1) is singular only at $r = 0$ [8]. The metric (1) is regular for $r > 0$ although this is manifest only on specification of D due to the compact form given. In particular, note that

$$g_{UU}|_{r=2m} = 2(D-2)(D-3)w^2. \tag{5}$$

Explicit forms of g_{UU} follow from (1) and for the cases $D = 4...11$ they are given in Appendix A. With U and w extending over the reals, (1) is geodesically complete. For the case $D = 4$ the coordinates are due to Israel [9].

For $D = 4$ a regular covering of the vacuum manifold is usually given in terms of the Kruskal-Szekeres coordinates (say u and v) [10] which, as is well known, are only implicit since $r(uv)$ is a non-invertible function. The generalization of these coordinates to D dimensions has been considered by Gregory and Laflamme [11]. (A. Krasiński has kindly pointed out an earlier study [12].) Now r is defined by way of a generalized tortoise coordinate so that

$$\pm\, 2muv = exp(\frac{D-3}{2m}\int\frac{r^{D-3}dr}{r^{D-3}-(2m)^{D-3}}) \equiv f(r)\frac{r-2m}{r+2m}. \tag{6}$$

Relatively simple forms for $f(r)$ are given in Appendix B for $D = 4...11$ excepting the cases $D = 8, 10$ for which no simple forms where found. Clearly the explicit form (2) is simpler. The disadvantage of the coordinates U, w lies in the fact that the "other" radial null geodesics are given in terms of the differential equation $dU/dw = -8m/g_{UU}$. Compared to the implicit nature of the Kruskal-Szekeres coordinates this disadvantage might be considered minor in the sense that these geodesics can always be constructed numerically. (Explicit solutions can be found depending on D.) Note that for all D, $dU/dw \to 0$ as $r \to 0$ and $dw/dU \to 0$ as $w \to 0$. The trajectories with D-tangents $l^a = \delta_U^a$ have $r\nabla_a l^a = (D-2)(D-3)w$ and become null geodesics at $w = 0$ where they generate the other branch to the horizon $r = 2m$.

Acknowledgments

This work was motivated by lectures at Black Holes IV and supported by a grant from the Natural Sciences and Engineering Research Council of Canada. Portions of this work were made possible by use of *GRTensorII* [13].

APPENDIX A: FORMS OF g_{UU} FOR $D = 4...11$

D=4 (Israel coordinates)

$$8\,\frac{mw^2}{r} \tag{A1}$$

D=5

$$16\,\frac{w^2(m+r)m}{r^2} \tag{A2}$$

D=6

$$8\,\frac{w^2(4\,m^2+4\,mr+3\,r^2)m}{r^3} \tag{A3}$$

D=7

$$16\,\frac{w^2(4\,m^3+4\,rm^2+3\,mr^2+2\,r^3)m}{r^4} \tag{A4}$$

D=8

$$8\,\frac{w^2(16\,m^4+16\,m^3r+12\,r^2m^2+8\,mr^3+5\,r^4)m}{r^5} \tag{A5}$$

D=9

$$16\,\frac{w^2(16\,m^5+16\,rm^4+12\,m^3r^2+8\,r^3m^2+5\,mr^4+3\,r^5)m}{r^6} \tag{A6}$$

D=10

$$8\,\frac{w^2(64\,m^6+64\,rm^5+48\,r^2m^4+32\,m^3r^3+20\,r^4m^2+12\,mr^5+7\,r^6)m}{r^7} \tag{A7}$$

D=11

$$16\,\frac{w^2(64\,m^7+64\,rm^6+48\,r^2m^5+32\,r^3m^4+20\,m^3r^4+12\,r^5m^2+7\,mr^6+4\,r^7)m}{r^8} \tag{A8}$$

APPENDIX B: FORMS FOR $\pm 2muv = f(r)\frac{r-2m}{r+2m}$

D=4 (Kruskal-Szekeres coordinates)

$$(r+2\,m)e^{\frac{r}{2m}} \tag{B1}$$

D=5

$$e^{\frac{r}{m}} \tag{B2}$$

D=6

$$\frac{(r+2\,m)e^{(\frac{3\,r}{2m}-\sqrt{3}\arctan(\frac{(r+m)}{\sqrt{3}m}))}}{\sqrt{r^2+2\,rm+4\,m^2}} \tag{B3}$$

D=7

$$e^{2\,(\frac{r}{m}-\arctan(\frac{r}{2m}))} \tag{B4}$$

D=9

$$e^{(\frac{3\,r}{m}-\sqrt{3}\arctan(\frac{(r+m)}{\sqrt{3}m})-\sqrt{3}\arctan(\frac{(r-m)}{\sqrt{3}m}))}(\frac{r^2-2\,mr\sqrt{2}+4\,m^2}{r^2+2\,mr\sqrt{2}+4\,m^2})^{\frac{1}{2}} \tag{B5}$$

D=11

$$e^{(\frac{4r}{m}-2\arctan(\frac{r}{2m})-\sqrt{2}\arctan(\frac{\sqrt{2}r-2\,m}{2m})-\sqrt{2}\arctan(\frac{\sqrt{2}r+2\,m}{2m}))}(\frac{r^2-2\,mr\sqrt{2}+4\,m^2}{r^2+2\,mr\sqrt{2}+4\,m^2})^{\frac{1}{\sqrt{2}}} \tag{B6}$$

---

[*] Electronic Address: lake@astro.queensu.ca

[1] R. Emparan and H. S. Reall, Phys. Rev. Lett. **88**, 101101 (2002) (hep-th/0110260).

[2] R. C. Myers and M. J. Perry, Annals Phys. **172**, 304 (1986) .

[3] S. Hwang, Geometriae Dedicata **71**, 5 (1998), G. W. Gibbons, D. Ida and T. Shiromizu, Prog. Theor. Phys. Suppl. **148**, 284 (2003) (gr-qc/0203004).

[4] F. R. Tangherlini, Nuovo Cimento **27**, 636 (1963).

[5] There are several recent studies concerning stability in higher dimensions. See, for example, G. W. Gibbons and S. A. Hartnoll, Phys. Rev. **D66**, 064024 (2002) (hep-th/0206202), G. W. Gibbons, S. A. Hartnoll and C. N. Pope, Phys. Rev. **D67**, 084024 (2003) (hep-th/0208031), V. Cardoso, O. J. C. Dias, J. P. S. Lemos, Phys. Rev. **D67**, 064026 (2003) (hep-th/0212168), A. Ishibashi and H. Kodama (hep-th/0305185), D. Birmingham (hep-th/0306004).

[6] The corresponding plane and hyperbollic cases are handled similarly.

[7] We use geometrical units. Note that the tensors here are $D \geq 4$ dimensional.

[8] Under a conformal transformation $\widetilde{g}_{ab} = \Omega^2 g_{ab}$, $\Omega = constant$, $\widetilde{R}_{ab} = R_{ab}$ but $\Omega^4 \widetilde{C}_{abcd}\widetilde{C}^{abcd} = C_{abcd}C^{abcd}$ and so the constant of proportionality in (4) is of no interest.

[9] W. Israel, Phys. Rev. **143**, 1016 (1966). The $D = 4$ Israel coordinates have been obtained independetly by D. W. Pajerski and E. T. Newman, J. Math. Phys. **12**, 1929 (1971) and by T. Klösch and T. Strobl, Class. Quant. Grav. **13**, 1191 (1996) (gr-qc/9507011).

[10] M. D. Kruskal, Phys. Rev. **119**, 1743 (1960), G. Szekeres, Publ. Math. Debrecen **7**, 285 (1960) (now available at General Relativity and Gravitation **34**, 2001, (2002)) .

[11] R. Gregory and R. Laflamme, Nuc. Phys. **B428**, 399 (1994) (hep-th/9404071).

[12] A. Krasiński and J. Plebański, Rep. Math. Phys. **17**, 217 (1980).

[13] This is a package which runs within Maple. It is entirely distinct from packages distributed with Maple and must be obtained independently. The GRTensorII software and documentation is distributed freely on the World-Wide-Web from the address *http://grtensor.org*